

VAXXON® ND-FLU

Protection right from the start



- Effective protection against Newcastle disease and Avian Influenza, strain H9N2
- Convenient, easy and safe to use

va:ino^{va}®

Protection right from the start

Newcastle disease (ND) is caused by virulent strains of avian paramyxovirus type 1 (APMV-1) of the genus Avulavirus which belongs to the family Paramyxoviridae. The ND virus (NDV) has shown to be able to infect over 200 species of birds, but the severity of the disease varies with both the host and the strain of the virus. Even APMV-1 strains of low virulence may induce severe respiratory diseases when exacerbated by the presence of other organisms or by adverse environmental conditions. Since its recognition in 1926, ND is regarded as being endemic in many countries (www.oie.int).

Avian influenza (AI) is a highly contagious viral disease affecting several species of food producing birds (chickens, turkeys, quails, guinea fowl), as well as pet birds and wild birds. Occasionally mammals, including humans, may contract Avian Influenza.

There are many AI virus strains, which are usually classified into two categories, according to the severity of the disease in poultry:

- low pathogenic (LPAI) strains, which typically cause few or no clinical signs in poultry.
- and highly pathogenic (HPAI) strains, which can cause severe clinical signs and potentially high mortality rates among poultry.

Avian Influenza has captured the attention of the international community over the years, with outbreaks in poultry having serious consequences on both livelihoods and international trade in many countries. In addition, although most avian influenza viruses do not infect humans, some, such as avian influenza H5N1 and H7N9, are well known to the public because of their implication in serious and sometimes fatal infections in people (www.oie.int).

H9N2 subtype avian influenza viruses (AIVs) have been isolated from various species of wild birds and domestic poultry in the world, and occasionally transmitted to humans. (1) In outbreaks of H9N2 AI reported in Iran, the isolate was characterized in the laboratory as of low pathogenicity (LP). However, mortality in the field approached 65% on some broiler farms. Lesions were identified only in the respiratory system, particularly the trachea and the bronchi. (2)

H9N2 remains endemic across Asia, mainly limited to outbreaks in domestic land-based poultry, but overshadowed as a pandemic threat by H5N1 bird flu, which has spread from Asia into Africa and Europe. However, there is evidence of interspecies transmission of H9N2 from land-based poultry to mammals, such as pigs and swine. Further evidence of an expanded mammalian host range includes efficient replication of H9N2 in mice without adaptation. H9N2 has already caused mild respiratory disease in humans in Hong Kong and mainland China in 1999 and 2003. (3)

Among low pathogenicity AIV (LPAIV), Eurasian H9N2 have been endemic in the domestic poultry population across North Africa, the Middle East and Asia since its emergence in China during 1994. These viruses are responsible for severe economic losses due to declined egg production and moderate to high mortality in broiler-type chickens. (4) In various outbreaks, inactivated influenza virus vaccines have been used successfully. (5)

Sources

- 1 - Dongdong Wang, Jingjing Wang, et All - Characterization of avian influenza H9N2 viruses isolated from ostriches (*Struthio camelus*) - Scientific Reports volume 8, Article number: 2273 [2018]
- 2 - H. Nili and K. Asasi - Avian Influenza (H9N2) Outbreak in Iran - Avian Diseases, 47(s3):828-831. 2003
- 3- Paul E Alexander, et all - Is H9N2 avian influenza virus a pandemic potential? - Can J Infect Dis Med Microbiol. 2009 Summer; 20(2): e35-e36.
- 4- Klaudia Chrzastek, et all - Characterization of H9N2 avian influenza viruses from the Middle East demonstrates heterogeneity at amino acid position 226 in the hemagglutinin and potential for transmission to mammals - Virology. 2018 May;518:195-201
- 5- Bahl, A. K., and B. S. Pomeroy. Efficacy of avian influenza oil-emulsion vaccine in breeder turkeys. J. Am. Vet. Med. Assoc. 171:1105. 1977

VAXXON® ND-FLU - Protection right from the start

broilers
layers
breeders

Target Species

Chicken (broilers, layers and breeders)

Indications

VAXXON® ND-FLU is for the active immunization against Newcastle disease and Avian Influenza A, subtype H9N2.

500 ds /
1000 ds

Dosage

Each vial contains 500 or 1000 doses.

Composition

Each 0.5 ml dose of vaccine contains sufficient:

- inactivated Newcastle disease virus.
- inactivated Avian Influenza A, strain H9N2.

day 1
onwards

Administration:

Broilers: 0.2 ml per chick at day old subcutaneously.

Layers & Breeders: 0.2 ml per bird from day old up to 3 weeks.

Re-vaccinations at 8 weeks and between 12-16 weeks are recommended with a dose of 0.5 ml intramuscularly.

24
months

Immunological properties

Stimulation of active immunity against Newcastle disease and Avian Influenza virus type A, subtype H9N2.

Shelf life

The product in the original packaging can be used until 24 months from the manufacturing date.

ND &
H9N2

Storage conditions

The product must be stored at 2°C to 8°C.

Do not freeze. Protect from light.

Disclaimer

This brochure may contain information on or may show or refer to products and/or services which are only available in certain countries. A reference to any product/service in this brochure does not imply that such product or such product with the same specifications et cetera, is or will be available at your location. Any product shown or referred to in this brochure may be subject to different regulatory requirements depending on the country of use. Please contact your local Vaxxinova representative for more information.

Vaxxinova® International BV does not give any warranty or other assurance as to the content of this brochure, its accuracy, completeness, timelessness or fitness for any particular purpose.

To the full extent permissible by law, Vaxxinova International BV disclaims all responsibility for any damages or losses (including without limitation, financial loss, damages for loss in business projects, loss of profits or other consequential losses) arising in contract, tort or otherwise from the use of inability to use any material of Vaxxinova, or from any action or decision taken as a result of using Vaxxinova's materials. Vaxxinova disclaims all responsibility and liability (including for negligence) in relation to information mentioned in Vaxxinova's materials.

This document or its content is not to be copied or further distributed without the consent of Vaxxinova International BV.

VAXXINOVA, VAXXON and GOVAXX are registered trademarks of Vaxxinova International B.V. (the Netherlands).

Version number 03, 2022.12 Product number 120855

Vaxxinova® International BV



Transistorweg 5
6534 AT Nijmegen, The Netherlands
+31 24 20 48 600
www.vaxxinova.com